

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 1 138 775 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
04.10.2001 Bulletin 2001/40

(51) Int Cl.7: **C12P 19/40**  
// (C12P19/40, C12R1:07),  
(C12P19/40, C12R1:19),  
(C12P19/40, C12R1:22)

(21) Application number: **01106745.1**

(22) Date of filing: **17.03.2001**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU**  
**MC NL PT SE TR**  
Designated Extension States:  
**AL LT LV MK RO SI**

Pub. No. 2001-269192

(30) Priority: **27.03.2000 JP 2000087302**

(71) Applicant: **YUKI GOSEI KOGYO CO., LTD.**  
**Chuo-ku Tokyo (JP)**

(72) Inventors:  
• **Mikami, Yoichi**  
**Tokyo (JP)**

• **Matsumoto, Seichiro**  
**Tokyo (JP)**  
• **Hayashi, Yoshinori**  
**Tokyo (JP)**  
• **Sato, Toyoki**  
**Tokyo (JP)**

(74) Representative:  
**Dreiss, Fuhlendorf, Steimle & Becker**  
**Patentanwälte,**  
**Postfach 10 37 62**  
**70032 Stuttgart (DE)**

(54) **Method of preparing a guanosine-group compound and an intermediate thereof**

(57) A glyoxal-guanosine-group compound is prepared either by reacting glyoxal-guanine with any one of ribose-1-phosphate and 2-deoxyribose-1-phosphate in the presence of purine nucleoside phosphorylase, or by reacting glyoxal-guanine with any one selected from the group consisting of uridine, 2'-deoxyuridine and thy-

midine, together with phosphate ion, in the presence of purine nucleoside phosphorylase and pyrimidine nucleoside phosphorylase. The glyoxal-guanosine-group compound is then decomposed by alkali, whereby a guanosine-group compound consisting of guanosine and 2'-deoxyguanosine is prepared.

EP 1 138 775 A2

## Description

[0001] The present invention relates to a method of preparing a guanosine-group compound, which is used as a raw material of an anti-virus agent, antisense medicine and the like, and a method of preparing a glyoxal-guanosine-group compound as an intermediate of the guanosine-group compound.


[0002] Guanosine or 2'-deoxyguanosine is industrially produced mainly by extraction/separation of hydrolysate of DNA (deoxyribonucleic acid) or RNA (ribonucleic acid), because the yield is extremely low when guanosine or 2'-deoxyguanosine is chemically synthesized. However, the hydrolysate of DNA contains 2'-deoxyadenosine, 2'-deoxycytidine and thymidine other than the targeted 2'-deoxyguanosine. The hydrolysate of RNA contains adenosine, cytidine and uridine other than the targeted guanosine. Accordingly, in order to collect only guanosine or 2'-deoxyguanosine, it is necessary to perform complicated processes of extraction/separation, thereby inevitably raising the production cost.

[0003] On the other hand, a method has been reported in which nucleoside (or deoxynucleoside) and nucleic acid base, which are the raw materials, are subjected to a base-exchange reaction by nucleoside phosphorylase, whereby the aimed nucleoside (or the aimed deoxynucleoside) is obtained (Hori, N., Watanabe, M., Yamazaki, Y., Mikami, Y., Agric. Biol. Chem., 53, 197-202 (1989)). By this method, adenosine and 2'-deoxyadenosine can be easily prepared (Jpn. Pat. Appln. KOKAI Publication No. 11-46790 "Method of preparing a purine nucleoside compound"). In order to obtain guanosine and 2'-deoxyguanosine by this method of enzymatic synthesis reaction, guanine must be soluble to water at least in a range of pH where the enzymes necessary for the reaction can function. In general, it is considered that the solubility of the reaction substrate is preferably at least equal to the concentration of Michaelis constant ( $K_m$ ) of the enzyme or higher, in order to carry out the enzyme reaction smoothly. However, as the solubility of guanine to water does not exceed a few ppm, the above method of enzymatic synthesis reaction cannot be employed in practice. Due to such an impasse-like situation, it has been demanded to develop a method of efficiently preparing guanosine or 2'-deoxyguanosine.

[0004] An object of the present invention is to provide a method of efficiently preparing guanosine or 2'-deoxyguanosine (i.e., a guanosine-group compound) at a high yield, as well as a method of efficiently preparing a glyoxal-guanosine or glyoxal-2'-deoxyguanosine (i.e., glyoxal-guanosine-group compound) as an intermediate of the guanosine-group compound at a high yield.


[0005] The inventors of the present invention have assiduously studied for solving the aforementioned problems. As a result, the inventors have discovered the following 1)-6).

1) When guanine, whose solubility to water does not exceed a few ppm, is reacted with glyoxal, glyoxal-guanine represented by the following formula (1) is obtained, and glyoxal-guanine is soluble to water.



(Glyoxal-guanine is also referred to as 6,7-dihydro-6,7-dihydroxyimidazo[1,2-a]purine-9(3H)-one.)

2) When microorganism itself which contains purine nucleoside phosphorylase (EC 2.4.2.1) or this enzyme derived from the microorganism are applied, for a reaction, to glyoxal-guanine and any one of ribose-1-phosphate and 2-deoxyribose-1-phosphate, a glyoxal-guanosine-group compound (i.e., glyoxal-guanosine or glyoxal-2'-deoxyguanosine) represented by the following formula (2) is obtained. Alternatively, when microorganism itself which contains purine nucleoside phosphorylase (EC 2.4.2.1) and pyrimidine nucleoside phosphorylase (EC 2.4.2.2) or these enzymes derived from the microorganism are applied, for a reaction, to glyoxal-guanine and any one selected from the group consisting of uridine, 2'-deoxyuridine and thymidine under the presence of phosphate ion, a glyoxal-guanosine-group compound (i.e., glyoxal-guanosine or glyoxal-2'-deoxyguanosine) represented by the following formula (2) is obtained. In other words, glyoxal-guanine can be a substrate of purine nucleoside phosphorylase.



(wherein R represents a hydrogen atom or a hydroxyl group.)

(Glyoxal-guanosine is also referred to as 3-( $\beta$ -D-erythropentofuranosyl)-6,7-dihydro-6,7-dihydroxyimidazo[1,2-a]purine-9(3H)-one, and glyoxal-2'-deoxyguanosine is also referred to as 3-(2-deoxy- $\beta$ -D-erythropentofuranosyl)-6,7-dihydro-6,7-dihydroxyimidazo[1,2-a]purine-9(3H)-one.)

3) In the method of preparing a glyoxal-guanosine-group compound described in 2) above, when at least one compound selected from the group consisting of glycine, iminodiacetic acid, nitrilotriacetic acid, ethylenediamine-tetraacetic acid (which will be referred to as "EDTA" hereinafter), ethylene glycol bis( $\beta$ -aminoethyl ether)-N,N,N',N'-tetraacetic acid (which will be referred to as "EGTA" hereinafter) and salts of these substances is added, or the above at least one compound is added in combination with boric acid or a salt thereof, the reaction yield is significantly improved. The above compound is added as a stabilizer of ribose-1-phosphate or 2-deoxyribose-1-phosphate, which is the substrate or an intermediate produced during base-exchange reaction.

4) As a principle, nucleoside phosphorylase derived from various organism can be efficient as the enzyme to be used in the preparation method of 2) above. However, the enzyme derived from microorganism of *Bacillus* genus, *Escherichia* genus or *Klebsiella* genus is suitable, and the thermotolerant enzyme derived from *Bacillus stearothermophilus* JTS 859 (FERM BP-6885) is especially preferable.


5) When an alkali is applied, for a reaction, to the glyoxal-guanosine-group compound obtained by the method of 2) above, the glyoxal portion of the glyoxal-guanosine-group compound is easily released, whereby a guanosine-group compound (i.e., guanosine or 2'-deoxyguanosine) can be obtained.

6) The preparation processes of the aforementioned 1)-5) can be carried out by one-pot reaction.


[0006] On the basis of the aforementioned discoveries, the present invention has been completed.

[0007] In summary, the present invention relates to the following methods (1)-(16).

(1) A method of preparing a glyoxal-guanosine-group compound represented by the formula (2):




wherein R represents a hydrogen atom or a hydroxyl group, which comprises the step of:  
reacting glyoxal-guanine represented by the formula (1):




10 with ribose-1-phosphate or 2-deoxyribose-1-phosphate in the presence of purine nucleoside phosphorylase, thereby obtaining a glyoxal-guanosine or glyoxal-2'-deoxyguanosine.

(2) A method of preparing a glyoxal-guanosine-group compound represented by the formula (2):




wherein R represents a hydrogen atom or a hydroxyl group, which comprises the step of:  
reacting glyoxal-guanine represented by the formula (1):




40 with any one selected from the group consisting of uridine, 2'-deoxyuridine and thymidine, together with phosphate ion, in the presence of purine nucleoside phosphorylase and pyrimidine nucleoside phosphorylase, thereby obtaining a glyoxal-guanosine or glyoxal-2'-deoxyguanosine.

(3) A method of preparing a guanosine-group compound, which comprises the steps of:  
reacting glyoxal-guanine represented by the formula (1):




55 with ribose-1-phosphate or 2-deoxyribose-1-phosphate in the presence of purine nucleoside phosphorylase, thereby obtaining a compound represented by the formula (2):




15 wherein R represents a hydrogen atom or a hydroxyl group; and  
decomposing, by alkali, the compound represented by the formula (2), thereby obtaining guanosine or 2'-  
deoxyguanosine.

(4) A method of preparing a guanosine-group compound, which comprises the steps of:  
reacting glyoxal-guanine represented by the formula (1):



30 with any one selected from the group consisting of uridine, 2'-deoxyuridine and thymidine, together with phosphate  
ion, in the presence of purine nucleoside phosphorylase and pyrimidine nucleoside phosphorylase, thereby ob-  
taining a compound represented by the formula (2):



50 wherein R represents a hydrogen atom or a hydroxyl group; and  
decomposing, by alkali, the compound represented by the formula (2), thereby obtaining guanosine or 2'-  
deoxyguanosine.

(5) A method of preparing a glyoxal-guanosine-group compound described in the aforementioned (1), wherein, as  
purine nucleoside phosphorylase, a microorganism itself which contains the enzyme or the enzyme derived from  
the microorganism is used.

55 (6) A method of preparing a glyoxal-guanosine-group compound described in the aforementioned (2), wherein, as  
purine nucleoside phosphorylase and pyrimidine nucleoside phosphorylase, a microorganism itself which contains  
the enzymes or the enzymes derived from the microorganism are used.

(7) A method of preparing a guanosine-group compound described in the aforementioned (3), wherein, as purine

nucleoside phosphorylase, a microorganism itself which contains the enzyme or the enzyme derived from the microorganism is used.

(8) A method of preparing a guanosine-group compound described in the aforementioned (4), wherein, as purine nucleoside phosphorylase and pyrimidine nucleoside phosphorylase, a microorganism itself which contains the enzymes or the enzymes derived from the microorganism are used.

(9) A method of preparing a glyoxal-guanosine-group compound described in the aforementioned (5) or (6), wherein the microorganism belongs to *Bacillus* genus, *Escherichia* genus or *Klebsiella* genus.

(10) A method of preparing a guanosine-group compound described in the aforementioned (7) or (8), wherein the microorganism belongs to *Bacillus* genus, *Escherichia* genus or *Klebsiella* genus.

(11) A method of preparing a glyoxal-guanosine-group compound described in any one of the aforementioned (5), (6) and (9), wherein the microorganism is *Bacillus stearothermophilus* JTS 859 (FERM BP-6885), *Escherichia coli* IFO 3301, *Escherichia coli* IFO 13168, or *Klebsiella pneumoniae* IFO 3321.

(12) A method of preparing a guanosine-group compound described in any one of the aforementioned (7), (8) and (10), wherein the microorganism is *Bacillus stearothermophilus* JTS 859 (FERM BP-6885), *Escherichia coli* IFO 3301, *Escherichia coli* IFO 13168, or *Klebsiella pneumoniae* IFO 3321.

(13) A method of preparing a glyoxal-guanosine-group compound described in any one of the aforementioned (1), (2), (5), (6), (9) and (11), wherein at least one compound selected from the group consisting of glycine, iminodiacetic acid, nitrilotriacetic acid, ethylenediaminetetraacetic acid, ethylene glycol bis ( $\beta$ -aminoethyl ether)-N,N,N',N'-tetraacetic acid and salts thereof is added, or the above at least one compound is added in combination with boric acid or a salt thereof.

(14) A method of preparing a guanosine-group compound described in any one of the aforementioned (3), (4), (7), (8), (10) and (12), wherein at least one compound selected from the group consisting of glycine, iminodiacetic acid, nitrilotriacetic acid, ethylenediaminetetraacetic acid, ethylene glycol bis( $\beta$ -aminoethyl ether)-N,N,N',N'-tetraacetic acid and salts thereof is added, or the above at least one compound is added in combination with boric acid or a salt thereof.

(15) A method of preparing a glyoxal-guanosine-group compound described in any one of the aforementioned (1), (2), (5), (6), (9), (11) and (13), which is performed by one-pot reaction.

(16) A method of preparing a guanosine-group compound described in any one of the aforementioned (3), (4), (7), (8), (10), (12) and (14), which is performed by one-pot reaction.

[0008] This summary of the invention does not necessarily describe all necessary features so that the invention may also be a sub-combination of these described features.

[0009] The present invention will be described in detail hereinafter.

#### <Raw material>

[0010] Glyoxal-guanine (i.e., 6,7-dihydro-6,7-dihydroxyimidazo[1,2-a]purine-9(3H)-one) to be used as a raw material in the present invention can be prepared at a high yield by adding guanine to a glyoxal aqueous solution and stirring the mixture with heating at a high temperature (preferably at 50-75°C) for about 15-20 hours. The solubility of glyoxal-guanine to water increases as the temperature rises. After the completion of the reaction, glyoxal-guanine is crystallized by cooling and can be obtained as a white solid by filtering.

[0011] However, when one-pot reaction is carried out by continuously adding nucleoside phosphorylase and either ribose donor or 2-deoxyribose donor, the solution in which glyoxal-guanine has been obtained may be used, as it is, as a raw material solution.

[0012] In the present invention, ribose-1-phosphate and uridine used as the ribose donor, and 2-deoxyribose-1-phosphate, 2'-deoxyuridine and thymidine used as the 2-deoxyribose donor are commercially available by, for example, Sigma Aldrich Japan Co.,.

#### <Synthesis conditions of a glyoxal-guanosine-group compound by the enzyme reaction>


[0013] The ribose donor or the 2-deoxyribose donor (which will be referred to as "the ribose-group donor" hereinafter) and glyoxal-guanine are added to a phosphate buffer solution (pH 6-8, preferably pH 7). Microorganism itself or the enzymes derived from the microorganism, which produces purine nucleoside phosphorylase and pyrimidine nucleoside phosphorylase (or only purine nucleoside phosphorylase), is added to the above phosphate buffer mixture. The resultant reaction mixture is stirred at the optimum temperature for the enzyme for 24-60 hours, resulting in reactions as represented by "Reaction formula 1 and Reaction formula 2" or "Reaction formula 3 and Reaction formula 4". Glyoxal-2'-deoxyguanosine is obtained by "Reaction formula 1 and Reaction formula 2". Glyoxal-guanosine is obtained by "Reaction formula 3 and Reaction formula 4". In Reaction formula 1 and Reaction formula 3 as the former half part,

pyrimidine nucleoside phosphorylase is used as the enzyme in the reaction. In Reaction formula 2 and Reaction formula 4 as the latter half part, purine nucleoside phosphorylase is used as the enzyme in the reaction. With respect to the optimum temperature for the enzyme, the temperature may be in the range of approximately 40-70°C in the case of the enzyme derived from *Bacillus stearothermophilus* JTS 859 (FERM BP-6885).


[0014] In the enzyme reaction of the present invention, purine nucleoside phosphorylase and pyrimidine nucleoside phosphorylase (or only purine nucleoside phosphorylase) are used. When pyrimidine nucleoside (uridine, 2'-deoxyuridine, or thymidine) is used as the ribose-group donor, pyrimidine nucleoside phosphorylase is essentially required for producing ribose-1-phosphate or 2-deoxyribose-1-phosphate in "Reaction formula 1" or "Reaction formula 3" as the former half reaction. On the other hand, when ribose-1-phosphate or 2-deoxyribose-1-phosphate is used as the ribose-group donor from the start of the reaction, the reaction represented by "Reaction formula 1" or "Reaction formula 3" is not necessitated, and thus pyrimidine nucleoside phosphorylase is not required. However, in this case, even if pyrimidine nucleoside phosphorylase is present, the overall enzyme reaction will not be adversely affected at all.

[0015] In Reaction formulae 1-4, "PYNP" represents pyrimidine nucleoside phosphorylase, and "PUNP" represents purine nucleoside phosphorylase.


(Reaction formula 1)



(Reaction formula 2)




(Reaction formula 3)





## (Reaction formula 4)



[0016] In the present invention, the initial concentration of the ribose-group donor as the raw material is 5-1000 mM, preferably 10-100 mM. Glyoxal-guanine does not dissolve in a large amount at a time. However, if it is assumed that the initially added amount of glyoxal-guanine were to be completely dissolved at a time, the expressed initial concentration would be 5-1000 mM, preferably 10-100 mM. The initial concentration of phosphate ion of 1 - 20 mM suffices the purpose.

<Stabilizer of deoxyribose-1-phosphate, etc.>

[0017] In the present invention, even when a nucleoside (uridine, 2'-deoxyuridine, thymidine) is used as the raw material, the reaction proceeds by way of ribose-1-phosphate or 2-deoxyribose-1-phosphate as the reaction intermediate, as shown in the aforementioned formulae. These phosphate compounds are unstable by nature and, when being left, naturally decomposed into ribose and phosphate ion or 2-deoxyribose and phosphate ion as time passes. When phosphatase is present in the reaction system, the decomposition is accelerated. If such decomposition occurs, the production efficiency of glyoxal-guanosine-group compound is decreased, because ribose or 2-deoxyribose, which is the product by decomposition of the intermediate, does not serve as the substrate of nucleoside phosphorylase. Therefore, in order to increase the production efficiency, a substance, which stabilizes ribose-1-phosphate or 2-deoxyribose-1-phosphate and inhibits the activity of phosphatase, has been searched in the present invention.

[0018] As a result, it has been discovered that addition of at least one compound selected from the group consisting of glycine, iminodiacetic acid, nitrilotriacetic acid, EDTA, EGTA and salts thereof, and more preferably, addition of the above at least one compound in combination with boric acid or a salt thereof is effective. The combination of EDTA or a salt thereof with boric acid or a salt thereof exhibits a significantly excellent effect. The preferable concentration of each composition added is 20-200 mM for boric acid or a salt thereof, and 2-10 mM for the above at least one compound or salts thereof (i.e., glycine, iminodiacetic acid, nitrilotriacetic acid, EDTA, EGTA). Salts of the above compounds and a salt of boric acid used herein are not particularly limited, and any salts can be used. For example, the salts used herein may be salts of alkaline metal such as sodium and potassium, or salts of alkaline earth metal such as calcium and magnesium.

## &lt;Preparation of a guanosine-group compound from a glyoxal-guanosine-group compound&gt;

[0019] A glyoxal-guanosine-group compound releases, by decomposition, the glyoxal portion thereof in an alkali aqueous solution, thereby obtaining a guanosine-group compound. As the alkali aqueous solution, 0.05-0.5N, preferably 0.1-0.25N sodium hydroxide aqueous solution can be used, for example. This decomposition reaction is accelerated by the increase in pH and temperature. However, it is essential to select a mild condition in which the bonding with ribose-group is not broken. Specifically, a glyoxal-guanosine-group compound is relatively stable in the range of pH 4-8, but is decomposed and releases the glyoxal portion thereof at pH 9 or higher. In conclusion, it is preferable that a glyoxal-guanosine-group compound is decomposed at 60-80°C in the range of pH 9-11 in 1-5 hours duration, to obtain a guanosine-group compound.

## &lt;Separation and purification of a guanosine-group compound&gt;

[0020] Collection of the product from the reaction solution can be performed by ultrafiltration, ion exchange separation, adsorption chromatography, crystallization and the like. The amount of the reaction product can be determined by the HPLC method using a UV detector.

## &lt;Microorganism and enzyme&gt;

[0021] Purine nucleoside phosphorylase (EC 2.4.2.1) and pyrimidine nucleoside phosphorylase (EC 2.4.2.2) used in the present invention may have any origin as a principle. The enzymes produced by the microorganism described below are preferable. The microorganism to be used in the present invention is not particularly limited to such examples, as long as the microorganism produces the aforementioned enzymes in a significant amount.

[0022] However, as shown in the aforementioned Reaction formulae, the present reactions employ nucleoside or 2'-deoxynucleoside as the substrate and thus inevitably produce a phosphate compound as the intermediate. Accordingly, microorganism which shows a strong nucleosidase and/or phosphatase activity cannot be utilized.

[0023] Examples of the microorganism which suffices these conditions include those which belong to *Bacillus* genus, *Escherichia* genus or *Klebsiella* genus. More specifically, the examples include *Bacillus stearothermophilus* JTS 859 (FERM BP-6885), *Escherichia coli* IFO 3301, *Escherichia coli* IFO 13168, and *Klebsiella pneumoniae* IFO 3321.

[0024] In the present reaction, the higher the solubility and the dissolving rate of the raw material are, the higher yield and the higher reaction rate are obtained. Accordingly, a relatively high temperature during the reaction is advantageous. Due to this, use of thermophilic bacteria and heat-resistant enzymes is preferable. Among the aforementioned examples of microorganisms, *Bacillus stearothermophilus* JTS 859 is the most preferable.

[0025] "IFO" indicates that the microorganisms are preserved in Institute for Fermentation, Osaka (IFO), 2-17-85, Zyusohonmachi, Yodogawa-ku, Osaka-shi, 532-0024 JAPAN. The "IFO strains" are available to any person, if desired. *Bacillus stearothermophilus* JTS 859 is under international deposit (the deposit number thereof is FERM BP-6885) in Patent Microorganism Depository, National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology. The microorganism JTS 859 (FERM BP-6885) was deposited on October 20, 1987 pursuant to BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDURE.

[0026] These types of microorganisms are grown on the normal culture medium for bacteria, and show better growth and produce a larger amount of the aforementioned enzyme by adding tryptone, yeast extract, glucose and the like. Further, addition of a nucleic acid compound such as inosine to the culture medium is effective in enhancing the activity of the enzyme. The cultured bacteria itself may be utilized as the crude enzyme. Alternatively, enzyme is obtained from the cultured bacteria by the standard methods (such as destruction by supersonication or milling, centrifugation, ammonium sulfate-fractionation, and membrane separation), and the resultant crude enzyme may also be employed.

[0027] The results of studying the bacteriological characteristics of the deposited strain according to "Bergey's Manual of Determinative Bacteriology, Volume II" (1984) are as follows. The experiments were basically carried out by the method described in "Biseibutsu no Bunrui to Dotei (Classification and Determination of Microorganism)" by Takeji Hasegawa, edited version, Gakkai Shuppan center, 1985.

[0028] *Bacillus stearothermophilus* JTS 859 (FERM BP-6885)

## 1. Morphological characteristics

(1) Shape and size of the cell: Rod, 5.4-6.5  $\mu$ m  $\times$  0.7-0.9  $\mu$ m

(2) Spore: Ellipsoidal-shaped spore is formed. One spore in one cell, the spore being positioned at the end of the cell.

## 2. Characteristics in culture

(1) Bouillon liquid culture: culture at 62°C for 2 days

(2) Bouillon agar plate culture: White with slightly yellow tint, glossy, opaque, not piling up. The colony has a wavy circular shape.

(3) Bouillon agar slant culture: White with slightly yellow tint, glossy, opaque, not piling up. Moderate growth.

## 3. Biochemical characteristics

(1)	Gram's stain	positive
(2)	Anaerobic culture	no growth
(3)	Motility	peritrichous movement
(4)	Oxidase	positive
(5)	Catalase	positive
(6)	Liquefiability of gelatin	liquefiable
(7)	Litmus milk	coagulated
(8)	O-F Test	fermentation type
(9)	VP Test	negative
(10)	Gas generation from D-glucose	no generation
(11)	Production of Acid from D-glucose	produced
(12)	Production of Acid from L-arabinose	not produced
(13)	Production of Acid from D-mannitol	produced
(14)	Production of Acid from D-xylose	not produced
(15)	Growth in Sabouraud's medium	
	Slant	Growth was observed
	Liquid	Growth was observed
(16)	Growth under 0.001 % lysozyme	no growth
(17)	Growth under 0.02 % azide	no growth
(18)	Growth under 7 % NaCl	no growth (grew up to 2% level)
(19)	Hydrolysis of casein	hydrolyzable
(20)	Hydrolysis of starch	hydrolyzable
(21)	Egg-yolk Test	no growth
(22)	Production of dihydroxyacetone	not produced
(23)	Utilization of citric acid	negative
(24)	Production of indole	negative
(25)	Activity of urease	positive
(26)	Deamination of phenylalanine	negative
(27)	Activity of arginine dihydrolase	positive
(28)	Decomposition of tyrosine	negative
(29)	Production of levan	negative
(30)	Reduction of a nitrate	positive
(31)	Denitrification of sodium nitrate	negative
(32)	Production of hydrogen sulfide	positive
(33)	Utilization of inorganic nitrogen source with NO <sub>3</sub> as the only nitrogen source with NH <sub>4</sub> as the only nitrogen source	Growth was observed Growth was observed
(34)	GC content	47.3 %
(35)	Growth temperature Range Optimum Range	40-71°C 60-68°C
(36)	Growth pH	

(continued)

	Range	5.7-8.5
	Optimum Range	6.0-7.0

[0029] Judging from the aforementioned characteristics, the present strain can be determined to be *Bacillus stearothermophilus*.

<one-pot reaction>

[0030] The method of preparing a glyoxal-guanosine-group compound and the method of preparing a guanosine-group compound of the present invention can be both carried out by a one-pot reaction. Specifically, the initial reaction solution is first prepared by reacting guanosine with a glyoxal aqueous solution. The ribose-group donor (preferably together with a stabilizer), the enzyme solution required for the reaction, and optionally the phosphate ion are added, for the reaction, to the initial reaction solution, whereby a glyoxal-guanosine-group compound can be prepared by a one-pot reaction. In addition, by alkali-decomposing the reaction product, a guanosine-group compound can be prepared by a one-pot reaction.

Examples

[0031] The present invention will further be described in detail by following Production Examples, Experiment Example, and Examples.

#### Production Example 1 Synthesis of glyoxal-guanine

[0032] 22.65g of guanine (manufactured by Tokyo Kasei Kogyo Co.,) and 21.75g of a 40% glyoxal aqueous solution (manufactured by Tokyo Kasei Kogyo Co.,) were added to 950 mL of purified water, and the mixture was stirred with heating at 70°C for 18 hours. The mixture was then cooled to 4°C and filtered. The white solid obtained by the filtration was dried, thereby resulting in 30.7g of glyoxal-guanine. The measurement result of nuclear magnetic resonance (NMR) spectrum is as follows.

[0033] <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 200 MHz) δ 8.52 (bs, 1H), 7.70 (bs, 1H), 7.16 (d, 1H), 6.39 (d, 1H), 5.46 (d, 1H), 4.83 (d, 1H).

#### Production Example 2 Preparation of bacteria suspension which contain nucleoside phosphorylase

[0034] *Bacillus stearothermophilus* JTS 859 (FERM BP-6885), which produces purine nucleoside phosphorylase (which will be referred to as "PUNP" hereinafter) and pyrimidine nucleoside phosphorylase (which will be referred to as "PYNP" hereinafter), was cultured in accordance with the following procedure.

[0035] In culturing the bacteria, a culture medium of pH 6.2 made of 10g of bacto-tryptone, 5g of yeast extract, 3g of glucose, 3g of salt, 1g of inosine and 1L of water was used. 100 mL of the aforementioned culture medium was charged into a 500 mL Erlenmeyer flask, a platinum loop of the bacteria collected from slant culture was inoculated into the medium in the flask, and the bacteria in the flask were cultured while shaking gyratingly (200 rpm) at 65°C for 16 hours, thereby resulting in cultured solution. 1.2L of the same culture medium was charged into a 2L-culturing-tank, and 60 mL of the cultured solution was inoculated into the medium in the tank. Then, the bacteria in the tank were cultured, while two stirring vanes (the upper vane and the lower vane, each having diameter of 42 mm) were rotated at 650 rpm, for 6 hours at the air flow rate of 1 vvm, at the culture temperature of 65°C and within the pH range of 6.5-7.0 (stat). After completing the culture, the bacteria were obtained by centrifuging (10,000G, 4°C, 15 minutes). The bacteria as a whole were suspended in 40 mL of 10 mM potassium phosphate solution (pH 7), whereby enzyme solution containing PUNP and PYNP (which will be referred to as "the enzyme solution" hereinafter) was obtained.

[0036] The enzyme activity required for converting 1 μmol of a substrate into a product in one minute is defined as "1U". The enzyme activity of the enzyme solution was measured by the standard method at 60°C. 1 mL of the enzyme solution exhibited the PUNP activity of 66 U and the PYNP activity of 166 U.

#### Experiment Example 1 Solubility of guanine and glyoxal-guanine

[0037] An excess amount of each compound was suspended in water, stirred for 1 hour at each temperature and left for a while. The liquid layer was analyzed with HPLC equipped with a UV detector, under the conditions described

below. The results are shown in Table 1.

**[0038]** Conditions of the HPLC analysis:

Detector: UV (260 nm)

Column: Superiorex ODS (manufactured by Shiseido)

Eluent: 0.1M ammonium dihydrogenphosphate solution: methanol=97:3 (v/v)

Flow rate: 1 mL/min

Sample: 20  $\mu$  L

Table 1

Compound	Solubility at each temperature (mg/kg)			
	30°C	40°C	50°C	60°C
Guanine	2 or less	2 or less	2	2
Glyoxal-guanine	200	275	330	440

**Example 1** Synthesis of glyoxal-2'-deoxyguanosine by nucleoside phosphorylase

**[0039]** 5 mM phosphate buffer (pH 7) containing 30 mM 2'-deoxyuridine, 30 mM glyoxal-guanine, 100 mM boric acid, and 4 mM disodium ethylenediaminetetraacetate was prepared. 1 mL of the enzyme solution obtained in Production Example 2 was added to 99 mL of the above phosphate buffer mixture, so that the total amount of the resultant reaction mixture was 100 mL. The reaction mixture was kept at 50°C for 48 hours, allowing the reaction to proceed. The mixture was then filtered at 60°C and cooled to 4°C. The white crystal obtained by the filtration (the yield was 80%) was subjected to spectrometry. The results are as follows. As a result of comparison with the values described in the reference (Chung, F., Hecht, S.S., Carcinogenesis, 6, 1671-1673 (1985)), it was confirmed that the product was glyoxal-2'-deoxyguanosine.

**[0040]** UV (pH 7),  $\lambda$  max (nm) 248, 272 (shoulder) 1H-NMR (DMSO-d<sub>6</sub>+D<sub>2</sub>O, 200 MHz),  $\delta$  7.95 (s, 1H, 2-H), 6.12 (t, 1H, 1'-H), 5.50 (s, 1H, 7-H), 4.90 (s, 1H, 6-H), 4.35 (bs, 1H, 3'-H), 3.83 (bs, 1H, 4'-H), 3.54 (m, 2H, 5'-H), 2.52 (m, 1H, 2'-H), 2.28 (m, 2H, 2'-H).

**Example 2** Synthesis of 2'-deoxyguanosine from glyoxal-2'-deoxyguanosine

**[0041]** 0.1g of glyoxal-2'-deoxyguanosine obtained in Example 1 described above was added to 0.2 N sodium hydroxide solution, and stirred at 70°C for 2 hours. After neutralizing the mixture, an analysis was performed by using the HPLC method, under the conditions of Example 1, thereby confirming that 2'-deoxyguanosine was quantitatively obtained. In addition, the white crystal obtained by chromatography and crystallization was subjected to spectrometry, and it was confirmed from each spectrum that the obtained crystal was 2'-deoxyguanosine.

**Example 3** The effect of the stabilizer on amount of glyoxal-2'-deoxyguanosine formed

**[0042]** 5 mM phosphate buffer (pH 7) containing 20 mM 2'-deoxyuridine, 20 mM glyoxal-guanine, the stabilizer shown in Table 2 and 1 mL of the enzyme solution prepared in Production Example 2 was prepared (In Table 2, "" indicates that the concentration of the stabilizer is 4 mM and the absence of "" indicates that the concentration of the stabilizer is 100 mM). 100 mL of the above phosphate buffer mixture was kept at 50°C for 38 hours, allowing the reaction to proceed. Glyoxal-2'-deoxyguanosine contained in the reaction solution was decomposed by alkali, to quantitatively obtain 2'-deoxyguanosine. Then, the concentration of 2'-deoxyguanosine was determined. Further, control solution was prepared in a manner similar to that of preparing the above buffer mixture except for adding 100 mM tris (hydroxymethyl) aminomethane hydrochloric acid ("Tris") instead of the stabilizer. This solution will be referred to as "Tris (control)" hereinafter. From the analysis on the results, it was discovered that the combination of a complex-forming compound similar to EDTA with boric acid is effective for improving the reaction yield.

Table 2

Stabilizer	Amount of glyoxal-2'-deoxyguanosine formed (mM)
Tris (control)	5.4
Boric acid	4.0

Table 2 (continued)

Stabilizer	Amount of glyoxal-2'-deoxyguanosine formed (mM)
Glycine	6.0
Iminodiacetic acid *	8.8
Nitrilotriacetic acid *	5.4
EDTA *	9.5
EGTA *	8.0
Tris and EDTA *	9.8
Glycine and EDTA *	10.1
Boric acid and Tris	5.6
Boric acid and glycine	6.5
Boric acid and iminodiacetic acid *	12.4
Boric acid and nitrilotriacetic acid *	12.2
Boric acid and EDTA *	12.8
Boric acid and EGTA *	11.5

#### Example 4 Synthesis of glyoxal-2'-deoxyguanosine by the enzyme derived from each strain

**[0043]** 500 mL of the same culture medium as used in Production Example 2 was charged into each of three 2L Erlenmeyer flasks. *Escherichia coli* IFO 3301, *Escherichia coli* IFO 13168 and *Klebsiella pneumoniae* IFO 3321 was each inoculated into the medium in each of three flasks, by collecting a platinum loop of bacteria each from the stock slant culture. The above bacteria each were cultured while shaking the flasks (200 rpm) at 37°C for 16 hours, thereby resulting in culture solution. The bacteria were collected by centrifuging (10,000 G, 4°C, 15 minutes) the culture solution. The obtained bacteria was suspended in 10 mL of 10 mM potassium phosphate buffer (pH 7), thereby resulting in three sets of enzyme solutions.

**[0044]** Glyoxal-2'-deoxyguanosine was synthesized from 2'-deoxyuridine and glyoxal-guanine under the following conditions, by using any one of the aforementioned three sets of enzyme solutions and the *Bacillus stearothermophilus* JTS 859 enzyme solution obtained in Production Example 2. Specifically, 5 mM phosphate buffer (pH 7) containing 30 mM 2'-deoxyuridine, 30 mM glyoxal-guanine, 100 mM boric acid, and 4 mM disodium ethylenediaminetetraacetate was first prepared. Each (2 mL) of the above four sets of enzyme solutions (which contain bacteria corresponding to 100 mL of the above culture solution) was added to 98 mL of the above phosphate buffer mixture, respectively, resulting in four sets of reaction mixtures. Each reaction mixture was kept at each reaction temperature for 48 hours, allowing the reaction to proceed. The results are shown in Table 3.

Table 3

Strain	Reaction temperature (°C)	Yield of glyoxal-2'-deoxyguanosine (%)
<i>Bacillus stearothermophilus</i> JTS 859	50	79
<i>Escherichia coli</i> IFO 3301	30	17
<i>Escherichia coli</i> IFO 13168	30	40
<i>Klebsiella pneumoniae</i> IFO 3321	30	23

#### Example 5 Synthesis of a glyoxal-guanosine-group compound by using various ribose-group donor

**[0045]** 5 mM phosphate buffer (pH 7) containing each ribose-group donor as shown in Table 4, 20 mM glyoxal-guanine, 100 mM boric acid, and 4 mM disodium ethylenediaminetetraacetate was first prepared (the concentration of each ribose-group donor was 20 mM. However, "" in Table 4 indicates that the concentration was 10 mM). 0.25 mL of the enzyme solution obtained in Production Example 2 was added to a required amount of the above phosphate

buffer mixture, so that the total amount of the resultant reaction mixture was 10 mL. The reaction mixture was kept at 50°C for 48 hours, allowing the reaction to proceed. The results are shown in Table 4.

Table 4

Ribose-group donor	Nucleoside formed	Yield (%)
Ribose-1-phosphate *	Glyoxal-guanosine	78
2-Deoxyribose-1-phosphate *	Glyoxal-2'-deoxyguanosine	76
Uridine	Glyoxal-guanosine	61
2'-Deoxyuridine	Glyoxal-2'-deoxyguanosine	79
Thymidine	Glyoxal-2'-deoxyguanosine	63

**Example 6** Effect of the substrate concentration on the preparation using the enzyme derived from *Bacillus stearothermophilus* JTS 859

**[0046]** 5 mM phosphate buffer (pH 7) containing guanine or glyoxal-guanine as the substrate base of each concentration shown in Table 5, 2'-deoxyuridine as the ribose-group donor having the same concentration as each substrate base, 100 mM boric acid, and 4 mM disodium ethylenediaminetetraacetate was first prepared. 1 mL of the enzyme solution obtained in Production Example 2 was added to 99 mL of the above phosphate buffer mixture. The resultant reaction mixture was kept at 50°C for 48 hours, allowing the reaction to proceed. The results are shown in Table 5. When glyoxal-guanine was selected as the substrate base, the formed glyoxal-2'-deoxyguanosine was decomposed by alkali, to obtain 2'-deoxyguanosine. The amount of the obtained 2'-deoxyguanosine was determined. In Table 5, the analysis was not performed when the concentration of guanine was no less than 50 mM.

Table 5

Concentration of substrate base (mM)	Yield of 2'-deoxyguanosine (%)						
	10 mM	20 mM	30 mM	50 mM	70 mM	80 mM	100 mM
Guanine (Comparative example)	29	23	19	-	-	-	-
Glyoxal-guanine	81	79	79	77	72	66	64

**Example 7** Synthesis by one-pot reaction

**[0047]** 0.45g of guanosine and 0.44g of a 40% glyoxal aqueous solution were added to 10 mL of water, and the mixture was stirred at 70°C for 18 hours. Next, the following substances and water were added to the above mixture, thereby resulting in the aqueous solution containing 30 mM 2'-deoxyuridine, 100 mM boric acid, 4 mM disodium ethylenediaminetetraacetate, and 5 mM potassium phosphate, as final concentration. 1 mL of the enzyme solution obtained in Production Example 2 was added to 99 mL of the above aqueous solution, so that the total volume of the resultant reaction mixture was 100 mL. The reaction mixture was kept at 50°C for 48 hours, allowing the reaction to proceed. Glyoxal-2'-deoxyguanosine contained in the reaction mixture was decomposed by alkali, to 2'-deoxyguanosine. Measurement of the concentration of 2'-deoxyguanosine by HPLC revealed that the yield was 80 %.

**Example 8** Synthesis of glyoxal-guanosine

**[0048]** 5 mM phosphate buffer (pH 7) containing 50 mM uridine, 50 mM glyoxal-guanine, 100 mM boric acid, and 4 mM disodium ethylenediaminetetraacetate was first prepared. 1 mL of the enzyme solution obtained in Production Example 2 was added to 99 mL of the above phosphate buffer mixture, so that the total volume of the resultant reaction mixture was 100 mL. The reaction mixture was kept at 50°C for 48 hours, allowing the reaction to proceed. After being filtered with heating, the mixture was cooled to 4°C. The white crystal obtained by the filtration (the yield was 72%) was subjected to spectrometry. The results are as follows. As a result of comparison with spectrum of glyoxal-2'-deoxyguanosine of Example 1, it was confirmed that the product was glyoxal-guanosine.

**[0049]** <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 200 MHz), δ 8.85 (s, 1H, 5-H), 8.00 (s, 1H, 2-H), 7.27 (d, 1H, 7-OH), 6.50 (d, 1H, 6-OH),


5.71 (d, 1H, 1'-H), 5.49 (d, 1H, 7-H), 5.44 (d, 1H, 3'-OH), 5.18 (d, 1H, 2'-OH), 5.06 (s, 1H, 5'-OH), 4.88 (d, 1H, 6-H), 4.41 (m, 1H, 2'-H), 4.10 (d, 1H, 3'-H), 3.90 (d, 1H, 4'-H), 3.57 (m, 2H, 5'-H).

[0050] Further, guanosine was prepared from the product obtained in this Example, in accordance with the procedure similar to that of Example 2. This fact also revealed that the product obtained in this Example was glyoxal-guanosine.


[0051] According to the present invention, a glyoxal-guanosine-group compound can be prepared from the ribose-group donor and the glyoxal-guanine, by the enzyme reactions derived from the microorganism. By alkali decomposing the glyoxal-guanosine-group compound, a guanosine-group compound consisting of guanosine and 2'-deoxyguanosine can be efficiently prepared at a high yield.

## Claims

1. A method of preparing a glyoxal-guanosine-group compound represented by the formula (2):




wherein R represents a hydrogen atom or a hydroxyl group, which comprises the step of;  
reacting glyoxal-guanine represented by the formula (1):




with ribose-1-phosphate or 2-deoxyribose-1-phosphate in the presence of purine nucleoside phosphorylase, thereby obtaining a glyoxal-guanosine or glyoxal-2'-deoxyguanosine.

2. A method of preparing a glyoxal-guanosine-group compound represented by the formula (2):






wherein R represents a hydrogen atom or a hydroxyl group, which comprises the step of:  
reacting glyoxal-guanine represented by the formula (1):



with any one selected from the group consisting of uridine, 2'-deoxyuridine and thymidine, together with phosphate ion, in the presence of purine nucleoside phosphorylase and pyrimidine nucleoside phosphorylase, thereby obtaining a glyoxal-guanosine or glyoxal-2'-deoxyguanosine.


15

3. A method of preparing a guanosine-group compound, which comprises the steps of:  
reacting glyoxal-guanine represented by the formula (1):



with ribose-1-phosphate or 2-deoxyribose-1-phosphate in the presence of purine nucleoside phosphorylase, thereby obtaining a compound represented by the formula (2):


30




wherein R represents a hydrogen atom or a hydroxyl group; and  
decomposing, by alkali, the compound represented by the formula (2), thereby obtaining guanosine or 2'-deoxyguanosine.

45

- 50 4. A method of preparing a guanosine-group compound, which comprises the steps of:  
reacting glyoxal-guanine represented by the formula (1):



10 with any one selected from the group consisting of uridine, 2'-deoxyuridine and thymidine, together with phosphate ion, in the presence of purine nucleoside phosphorylase and pyrimidine nucleoside phosphorylase, thereby obtaining a compound represented by the formula (2):




wherein R represents a hydrogen atom or a hydroxyl group; and

30 decomposing, by alkali, the compound represented by the formula (2), thereby obtaining guanosine or 2'-deoxyguanosine.

- 35
5. A method of preparing a glyoxal-guanosine-group compound according to claim 1, **characterized in that**, as purine nucleoside phosphorylase, a microorganism itself which contains said enzyme or said enzyme derived from the microorganism is used.
  - 40 6. A method of preparing a glyoxal-guanosine-group compound according to claim 2, **characterized in that**, as purine nucleoside phosphorylase and pyrimidine nucleoside phosphorylase, a microorganism itself which contains said enzymes or said enzymes derived from the microorganism are used.
  - 45 7. A method of preparing a guanosine-group compound according to claim 3, **characterized in that**, as purine nucleoside phosphorylase, a microorganism itself which contains said enzyme or said enzyme derived from the microorganism is used.
  8. A method of preparing a guanosine-group compound according to claim 4, **characterized in that**, as purine nucleoside phosphorylase and pyrimidine nucleoside phosphorylase, a microorganism itself which contains said enzymes or said enzymes derived from the microorganism are used.
  9. A method of preparing a glyoxal-guanosine-group compound according to claim 5 or 6, **characterized in that** the microorganism belongs to *Bacillus* genus, *Escherichia* genus or *Klebsiella* genus.
  - 50 10. A method of preparing a guanosine-group compound according to claim 7 or 8, **characterized in that** the microorganism belongs to *Bacillus* genus, *Escherichia* genus or *Klebsiella* genus.
  - 55 11. A method of preparing a glyoxal-guanosine-group compound according to any one of claims 5, 6 and 9, **characterized in that** the microorganism is *Bacillus stearothermophilus* JTS 859 (FERM BP-6885), *Escherichia coli* IFO 3301, *Escherichia coli* IFO 13168, or *Klebsiella pneumoniae* IFO 3321.
  12. A method of preparing a guanosine-group compound according to any one of claims 7, 8 and 10, **characterized**

In that the microorganism is *Bacillus stearothermophilus* JTS 859 (FERM BP-6885), *Escherichia coli* IFO 3301, *Escherichia coli* IFO 13168, or *Klebsiella pneumoniae* IFO 3321.

- 5 13. A method of preparing a glyoxal-guanosine-group compound according to any one of claims 1, 2, 5, 6, 9 and 11, characterized in that at least one compound selected from the group consisting of glycine, iminodiacetic acid, nitrilotriacetic acid, ethylenediaminetetraacetic acid, ethylene glycol bis (β-aminoethyl ether)-N,N,N',N'-tetraacetic acid and salts thereof is added, or the above at least one compound is added in combination with boric acid or a salt thereof.
- 10 14. A method of preparing a guanosine-group compound according to any one of claims 3, 4, 7, 8, 10 and 12, characterized in that at least one compound selected from the group consisting of glycine, iminodiacetic acid, nitrilotriacetic acid, ethylenediaminetetraacetic acid, ethylene glycol bis(β-aminoethyl ether)-N,N,N',N'-tetraacetic acid and salts thereof is added, or the above at least one compound is added in combination with boric acid or a salt thereof.
- 15 15. A method of preparing a glyoxal-guanosine-group compound according to any one of claims 1, 2, 5, 6, 9, 11 and 13, which is performed by one-pot reaction.
- 20 16. A method of preparing a guanosine-group compound according to any one of claims 3, 4, 7, 8, 10, 12 and 14, which is performed by one-pot reaction.



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 1 138 775 A3**

(12) **EUROPEAN PATENT APPLICATION**

(88) Date of publication A3:  
17.04.2002 Bulletin 2002/16

(43) Date of publication A2:  
04.10.2001 Bulletin 2001/40

(21) Application number: 01106745.1

(22) Date of filing: 17.03.2001

(51) Int Cl.7: **C12P 19/40**, C12P 19/18,  
C12P 19/28, C12N 1/20,  
C07H 1/00, C07H 19/167,  
C07H 19/173  
// (C12P19/40, C12R1:07),  
(C12P19/40, C12R1:19),  
(C12P19/40, C12R1:22)

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU**  
**MC NL PT SE TR**  
Designated Extension States:  
**AL LT LV MK RO SI**

(30) Priority: 27.03.2000 JP 2000087302

(71) Applicant: YUKI GOSEI KOGYO CO., LTD.  
Chuo-ku Tokyo (JP)

(72) Inventors:  
• Mikami, Yoichi  
Tokyo (JP)

• Matsumoto, Selchiro  
Tokyo (JP)  
• Hayashi, Yoshinori  
Tokyo (JP)  
• Sato, Toyoki  
Tokyo (JP)

(74) Representative:  
Dreiss, Fuhlendorf, Stelmle & Becker  
Patentanwälte,  
Postfach 10 37 62  
70032 Stuttgart (DE)

(54) **Method of preparing a guanosine-group compound and an intermediate thereof**

(57) A glyoxal-guanosine-group compound is prepared either by reacting glyoxal-guanine with any one of ribose-1-phosphate and 2-deoxyribose-1-phosphate in the presence of purine nucleoside phosphorylase, or by reacting glyoxal-guanine with any one selected from the group consisting of uridine, 2'-deoxyuridine and thymidine, together with phosphate ion, in the presence of

purine nucleoside phosphorylase and pyrimidine nucleoside phosphorylase. The glyoxal-guanosine-group compound is then decomposed by alkali, whereby a guanosine-group compound consisting of guanosine and 2'-deoxyguanosine is prepared.

EP 1 138 775 A3



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 01 10 6745

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Y,D	EP 0 896 065 A (YUKI GOSEI KOGYO CO., LTD.) 10 February 1999 (1999-02-10) see especially the Examples * the whole document *	1-16	C12P19/40 C12P19/18 C12P19/28 C12N1/20 C07H1/00
Y	EP 0 411 158 A (YAMASA SHOYU KABUSHIKI KAISHA) 6 February 1991 (1991-02-06) see e.g. Table 5 on p. 14 * the whole document *	1-16	C07H19/167 C07H19/173 //(C12P19/40, C12R1:07), (C12P19/40, C12R1:19), (C12P19/40, C12R1:22)
A	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; CHUNG, FUNG LUNG ET AL: "Formation of the cyclic 1,N2-glyoxal-deoxyguanosine adduct upon reaction of N-nitroso-2-hydroxymorpholine with deoxyguanosine" retrieved from STN Database accession no. 104:16371 XP002191262 * abstract * & CARCINOGENESIS (LONDON) (1985), 6(11), 1671-3 ,  -----	1-16	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			C12P C12N C07H
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 25 February 2002	Examiner Douschan, K
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			

EPO FORM 1503 (3.92) (P04001)

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 10 6745

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

25-02-2002

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
EP 0896065	A	10-02-1999	JP	11046790 A	23-02-1999
			CN	1207417 A	10-02-1999
			CZ	9802419 A3	17-02-1999
			EP	0896065 A1	10-02-1999
			US	6017736 A	25-01-2000
EP 0411158	A	06-02-1991	DE	69027154 D1	04-07-1996
			DE	69027154 T2	24-10-1996
			EP	0411158 A1	06-02-1991
			KR	9710134 B1	21-06-1997
			US	5506122 A	09-04-1996
			AT	138689 T	15-06-1996
			AU	631607 B2	03-12-1992
			AU	5108090 A	26-09-1990
			CA	2028119 A1	29-08-1990
			DK	411158 T3	17-06-1996
			ES	2089010 T3	01-10-1996
			WO	9010080 A1	07-09-1990
			JP	2795748 B2	10-09-1998
			US	5384251 A	24-01-1995
			US	5258301 A	02-11-1993

EPO FORM P449

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82